



NETTLEHAM JUNIOR SCHOOL – SICKNESS/ABSENCE REPORTING

If your child is unwell please ring school (or email us on enquiries@nettleham-junior.lincs.sch.uk) before 10.00 a.m. on the day your child's illness starts. You are also able to report a child's absence from school via our Website using this link: <https://www.nettleham-junior.lincs.sch.uk/page/?title=Sickness+Reporting&pid=131>.

We have a designated option on our new phone system where you are able to leave a message reporting an absence. However, it is essential that we know **exactly** what it is your child is unwell with – we have some messages simply saying their child is “poorly” or “sick”. In order for us to keep track of what illnesses we have in school it is essential that you let us have the precise nature of any sickness – **particularly if it is an infectious disease i.e. chickenpox, measles, german measles, mumps, whooping cough, scarlet fever etc. There are children in school who have medical conditions which can be seriously affected by contact with these illnesses.**

If your child is going to be absent for longer than 48 hours could you please report his/her absence again with an update. On your child's return please send a letter or email, outlining the details of the illness, confirming how long they have been absent and that they are fit to return to school.

We also get queries regarding how long a child should be kept away from school. We follow the guidelines issued by the Health Protection Agency a copy of which is attached. Should you require a paper copy please call into school.

Thank you for your co-operation in this matter and your continued support of our school.

Yours sincerely,

Mr. D. Gibbons
Headteacher



Rashes and skin infections	Recommended period to be kept away from school, nursery or childminders	Comments
Athlete's foot	None	Athlete's foot is not a serious condition. Treatment is recommended
Chickenpox*	Until all vesicles have crusted over	See: Vulnerable children and female staff – pregnancy
Cold sores, (Herpes simplex)	None	Avoid kissing and contact with the sores. Cold sores are generally mild and self-limiting
German measles (rubella)*	Four days from onset of rash (as per "Green Book")	Preventable by immunisation (MMR x 2 doses). See: Female staff – pregnancy
Hand, foot and mouth	None	Contact the Duty Room if a large number of children are affected. Exclusion may be considered in some circumstances
Impetigo	Until lesions are crusted and healed, or 48 hours after commencing antibiotic treatment	Antibiotic treatment speeds healing and reduces the infectious period
Measles*	Four days from onset of rash	Preventable by vaccination (MMR x 2). See: Vulnerable children and female staff – pregnancy
Molluscum contagiosum	None	A self-limiting condition
Ringworm	Exclusion not usually required	Treatment is required
Roseola (infantum)	None	None
Scabies	Child can return after first treatment	Household and close contacts require treatment
Scarlet fever*	Child can return 24 hours after commencing appropriate antibiotic treatment	Antibiotic treatment recommended for the affected child. If more than one child has scarlet fever contact PHA Duty Room for further advice
Slapped cheek (fifth disease or parvovirus B19)	None once rash has developed	See: Vulnerable children and female staff – pregnancy
Shingles	Exclude only if rash is weeping and cannot be covered	Can cause chickenpox in those who are not immune i.e. have not had chickenpox. It is spread by very close contact and touch. If further information is required, contact the Duty Room. SEE: Vulnerable Children and Female Staff – Pregnancy
Warts and verrucae	None	Verrucae should be covered in swimming pools, gymnasiums and changing rooms

Diarrhoea and vomiting illness	Recommended period to be kept away from school, nursery or childminders	Comments
Diarrhoea and/or vomiting	48 hours from last episode of diarrhoea or vomiting	
<i>E. coli</i> O157 VTEC*	Should be excluded for 48 hours from the last episode of diarrhoea	Further exclusion is required for young children under five and those who have difficulty in adhering to hygiene practices
Typhoid* [and paratyphoid*] (enteric fever)	Further exclusion may be required for some children until they are no longer excreting	Children in these categories should be excluded until there is evidence of microbiological clearance. This guidance may also apply to some contacts of cases who may require microbiological clearance
Shigella* (dysentery)		Please consult the Duty Room for further advice
Cryptosporidiosis*	Exclude for 48 hours from the last episode of diarrhoea	Exclusion from swimming is advisable for two weeks after the diarrhoea has settled

Respiratory infections	Recommended period to be kept away from school, nursery or childminders	Comments
Flu (influenza)	Until recovered	See: Vulnerable children
Tuberculosis*	Always consult the Duty Room	Requires prolonged close contact for spread
Whooping cough* (pertussis)	48 hours from commencing antibiotic treatment, or 21 days from onset of illness if no antibiotic treatment	Preventable by vaccination. After treatment, non-infectious coughing may continue for many weeks. The Duty Room will organise any contact tracing necessary

Other infections	Recommended period to be kept away from school, nursery or childminders	Comments
Conjunctivitis	None	If an outbreak/cluster occurs, consult the Duty Room
Diphtheria *	Exclusion is essential. Always consult with the Duty Room	Family contacts must be excluded until cleared to return by the Duty Room. Preventable by vaccination. The Duty Room will organise any contact tracing necessary
Glandular fever	None	
Head lice	None	Treatment is recommended only in cases where live lice have been seen
Hepatitis A*	Exclude until seven days after onset of jaundice (or seven days after symptom onset if no jaundice)	The duty room will advise on any vaccination or other control measure that are needed for close contacts of a single case of hepatitis A and for suspected outbreaks.
Hepatitis B*, C, HIV/AIDS	None	Hepatitis B and C and HIV are bloodborne viruses that are not infectious through casual contact. For cleaning of body fluid spills. SEE: Good Hygiene Practice
Meningococcal meningitis*/septicaemia*	Until recovered	Some forms of meningococcal disease are preventable by vaccination (see immunisation schedule). There is no reason to exclude siblings or other close contacts of a case. In case of an outbreak, it may be necessary to provide antibiotics with or without meningococcal vaccination to close contacts. The Duty Room will advise on any action needed.
Meningitis* due to other bacteria	Until recovered	Hib and pneumococcal meningitis are preventable by vaccination. There is no reason to exclude siblings or other close contacts of a case. The Duty Room will give advice on any action needed
Meningitis viral*	None	Milder illness. There is no reason to exclude siblings and other close contacts of a case. Contact tracing is not required
MRSA	None	Good hygiene, in particular handwashing and environmental cleaning, are important to minimise any danger of spread. If further information is required, contact the Duty Room
Mumps*	Exclude child for five days after onset of swelling	Preventable by vaccination (MMR x 2 doses)
Threadworms	None	Treatment is recommended for the child and household contacts
Tonsillitis	None	There are many causes, but most cases are due to viruses and do not need an antibiotic

* denotes a notifiable disease. It is a statutory requirement that doctors report a notifiable disease to the Director of Public Health via the Duty Room.

Outbreaks: if a school, nursery or childminder suspects an outbreak of infectious disease, they should inform the Duty Room.